



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SCIENCE

FRIDAY, DECEMBER 15, 1911

PRINCIPLES OF WATER-POWER DEVELOPMENT¹

CONTENTS

<i>Principles of Water-power Development:</i> DR. W J MCGEE	813
<i>University Extension and the State University:</i> PROFESSOR LOUIS E. REBER	825
<i>An Opportunity for the Spirit of Research in Laboratory Instruction in Physics:</i> PROFESSOR G. W. STEWART	833
<i>A New Gas Volcano in Trinidad:</i> ROBERT ANDERSON	834
<i>The Future of the London Zoological Gardens</i>	835
<i>Scientific Notes and News</i>	836
<i>University and Educational News</i>	840
<i>Discussion and Correspondence:—</i>	
“Genotype” and “Pure Line”: PROFESSOR H. S. JENNINGS. <i>Mosquito Romance:</i> KONOPS. <i>How a Falling Cat turns over in the Air, How to throw a Curved Ball:</i> PROFESSOR W. S. FRANKLIN. <i>Seed Distribution by Surface Tension:</i> HENRY S. CONARD. <i>Models of Vorticella and Cyclops:</i> PROFESSOR HENRY F. NACHTRIEB. <i>Siphon Springs and Sink Holes:</i> PROFESSOR H. F. CLELAND. <i>The Rôle of Salts in the Preservation of Life:</i> PROFESSOR JACQUES LOEB .	841
<i>Scientific Books:—</i>	
<i>Observations and Investigations made at the Blue Hill Observatory:</i> PROFESSOR R. DEC. WARD. <i>Iddings's Rock Minerals:</i> DR. GEO. P. MERRILL. <i>Alder and Hancock on British Nudibranchiate Mollusca; Antarctic Expedition of the Duke of Orleans:</i> DR. WM. H. DALL. <i>Newbiggin's Modern Geography:</i> PROFESSOR RICHARD W. DODGE	846
<i>The Convocation Week Meetings of Scientific Societies</i>	851

MSS. intended for publication and books, etc., intended for review should be sent to the Editor of SCIENCE, Garrison-on-Hudson, N. Y.

1. The development of water-power involves artificial regulation of streams. Proper regulation of running water for the several uses of water supply, irrigation, power and navigation can be effected only in the light of the physical relations, the relations in equity, and the more salient legal relations of water in streams.

PHYSICAL RELATIONS

2. The fresh water of the land is derived directly from rainfall (including snow) and indirectly through evaporation from the sea. The mean annual rainfall on mainland United States ranges from less than 5 to over 100 inches, averaging 30 inches; the quantity aggregates about 5,000,000,000 acre-feet.² The distribution is unequal; over the eastward two fifths of the country the mean is about 48 inches, over the median fifth some 30 inches, and over the westward two fifths about 12 inches.³

3. In humid lands the water of rains and melting snows tends to gather into streams, generally taking the shortest and easiest paths to the sea, while in arid lands

¹ Presented at a hearing of the National Waterways Commission, November 21, 1911.

² The acre-foot is a convenient unit not only because in common use throughout arid America, but because large enough to measure water in its national aspect without use of incomprehensibly large figures. It equals 43,560 cubic feet, 326,047 gallons, or 1,359.6 tons; it is something over a kilostere (equaling 1.2335 ks.), or cube of 10 meters.

³ “Soil Erosion,” Bureau of Soils Bulletin 71, 1911, p. 17.